

Dynamical renormalization group approach to the Altarelli-Parisi-Lipatov equations

D. Boyanovsky,¹ H. J. de Vega,² D.-S. Lee,³ S.-Y. Wang,¹ and H.-L. Yu⁴

¹*Department of Physics and Astronomy, University of Pittsburgh, Pittsburgh, Pennsylvania 15260, USA*

²*LPTHE, Université Pierre et Marie Curie (Paris VI) et Denis Diderot (Paris VII),*

Tour 16, 1er. étage, 4, Place Jussieu, 75252 Paris, Cedex 05, France

³*Department of Physics, National Dong Hwa University, Shoufeng, Hualien 974, Taiwan, ROC*

⁴*Institute of Physics, Academia Sinica, Taipei, Taiwan 115, ROC*

(Dated: February 1, 2008)

The Altarelli-Parisi-Lipatov equations for the parton distribution functions are rederived using the dynamical renormalization group approach to quantum kinetics. This method systematically treats the $\ln Q^2$ corrections that arises in perturbation theory as a renormalization of the parton distribution function and unambiguously indicates that the strong coupling must be allowed to run with the scale in the evolution kernel. To leading logarithmic accuracy the evolution equation is Markovian and the logarithmic divergences in the perturbative expansion are identified with the secular divergences (terms that grow in time) that emerge in a perturbative treatment of the kinetic equations in nonequilibrium systems. The resummation of the leading logarithms by the Altarelli-Parisi-Lipatov equation is thus similar to the resummation of the leading secular terms by the Boltzmann kinetic equation.

PACS numbers: 11.10.Hi, 12.38.Aw, 11.10.-z

An early important test of QCD and one that validates the parton picture, is the Q^2 evolution of the quark distribution, or structure function, in deep inelastic lepton-hadron scattering which reveals scaling violations as predicted by QCD. The leading logarithmic dependence of the quark structure function on the scale Q^2 in deeply inelastic scattering was originally computed in QCD via the operator product expansion [1]. In perturbation theory, parton emission leads to QCD corrections to the distribution functions that grow as $\ln Q^2$, which eventually become large and invalidate the perturbative expansion. These corrections signal the violations of scaling in deep inelastic scattering.

Lipatov [2] and Altarelli and Parisi [3] provided an alternative derivation of the evolution equations for the parton distribution functions by associating the $\ln Q^2$ corrections with a *change* in the parton distribution function at the scale Q^2 . These authors provided a set of integro-differential equations that resum the perturbative expansion and describe the evolution of the parton densities as a function of $\ln Q^2$ [4]. These parton evolution equations are successful in predicting the correct leading logarithmic dependence of the distribution functions in Q^2 and the scaling violations predicted by these equations have been confirmed spectacularly in deep inelastic experiments [5].

Altarelli and Parisi used a probabilistic description in their derivation, however infrared singularities associated with the emission of low momentum partons introduce singularities in some probability densities. Anticipating that the infrared singularities should cancel between real and virtual soft gluons, these authors provided a prescription to handle these singularities.

Collins and Qiu [6] provided a derivation of the Altarelli-Parisi-Lipatov (APL) equations with a special treatment of virtual diagrams that clearly reveals the explicit cancellation of the infrared singularities between the contributions of real and virtual soft gluons through Ward identities. The result of Collins and Qiu is contained in a set of integro-differential equations, equivalent to the APL ones but without the need of extra prescriptions for handling soft gluon emission since the cancellation of infrared singularities is manifest in their treatment of real and virtual diagrams. For valence quarks these equations have the form

$$Q^2 \frac{d}{dQ^2} q_i(x, Q^2) = \frac{\alpha_s(Q^2)}{2\pi} \left[\int_x^1 \frac{dx_1}{x_1} q_i(x_1, Q^2) \gamma_{qq} \left(\frac{x}{x_1} \right) - q_i(x, Q^2) \int_0^1 dz \gamma_{qq}(z) \right], \quad (1)$$

where $q_i(x, Q^2)$ is the distribution function of the (valence) i -quark at a fixed momentum fraction x that evolves with the momentum scale Q^2 at which the distribution function is probed and the quark-quark splitting function is given by

$$\gamma_{qq}(z) = \frac{N_c^2 - 1}{2N_c} \left(\frac{1+z^2}{1-z} \right) \quad (2)$$

with N_c the number of colors. The splitting function $\gamma_{qq}(z)$ describe the probability density of finding a parton quark inside another parton quark with momentum fraction z of the parent parton momentum and was originally

introduced by Weiszacker and Williams for QED [7]. The relative minus sign between the two terms allows an explicit cancellation of the infrared divergences arising from (real and virtual) soft gluon emission [6].

A remarkable aspect of the Altarelli-Parisi-Lipatov equations in the form obtained by Collins and Qiu given by Eq. (1) above, is that they feature exactly the same splitting function in both terms on the right-hand side. This feature allows an unambiguous probabilistic interpretation as a set of *Boltzmann kinetic equations*, since the different contributions can be interpreted as “gain” and “loss” processes that change the parton distribution in “cells” labelled by the variable x , the fractional momentum of the hadron carried by the parton [6, 8].

Within the context of the probabilistic interpretation suggested by Collins and Qiu and for reasons that will become clear below, it is important to remark at this stage that the Altarelli-Parisi-Lipatov equation involves a convolution in the variable x but *not* in the variable Q^2 , i.e., the parton distribution function on the right-hand side of the equation is evaluated at the same Q^2 as that on the left-hand side. This feature of the evolution equation results in a *Markovian* approximation of parton evolution inside a hadron. Physically, this implies that the evolution of the parton distribution at some fixed x involves only parton emission or absorption processes taking place at *momentum scale* Q^2 where the parton distribution is probed, rather than those taking place at any other momentum scales. This property is actually inherent from the leading logarithmic approximation employed in the formulation, where each parton emission or absorption process is actually independent of the others and depends only on the momentum scale at which the process takes place.

This interpretation of the Altarelli-Parisi-Lipatov equations as a set of kinetic equations for the parton distribution functions motivates us to provide yet another alternative derivation of these equations by implementing a novel approach to quantum kinetics introduced recently in the form of the *dynamical renormalization group* [9].

While we anticipate that the approach presented here will ultimately lead to the form of the Altarelli-Parisi-Lipatov equations derived by Collins and Qiu, we believe that this alternative formulation yields novel insights that could prove valuable to implement a systematic treatment of parton evolution. As will be shown in detail below, the rederivation of the evolution equations via the dynamical renormalization group provides further and deeper understanding of many important aspects:

- (i) It reveals some subtle aspects of the derivation which highlight the role of the running of the strong coupling and the “separation of scales” which is a necessary ingredient in a kinetic description. The role of the running coupling is ultimately at the heart of the scaling violations and the necessity for including the running of the coupling in the evolution kernel becomes manifest in this approach.
- (ii) The dynamical renormalization group directly reveals how the $\ln Q^2$ that arise in the perturbative expansion can be systematically absorbed into the parton distribution functions and establishes a direct correspondence with *secular divergences in time* [9] that are manifest in the derivation of the Boltzmann equation and also responsible for the failure of the perturbative expansion in quantum kinetics.
- (iii) It directly establishes a correspondence between the leading logarithmic approximation to parton evolution equation which is local in Q^2 and the Markovian approximation to kinetic equation which is local in time.

As noticed by Collins and Qiu and mentioned above, the connection between the Altarelli-Parisi-Lipatov equation and the kinetic equation in nonequilibrium statistical physics is established by identifying $\ln Q^2$ with the “time” variable. This identification is the main concept upon which the formulation based on the dynamical renormalization group hinges. In this article, we revisit the derivation of Eq. (1) but implementing the concepts of the dynamical renormalization group [9].

The i -quark distribution function at some scale Q^2 with fixed momentum fraction x is defined as [6]

$$q_i(x, Q^2) = (2\pi)^3 2E_p \int \frac{d^4 p}{2P \cdot n} T_{ba}^i(P, p) (\gamma_-)_{ba} \delta\left(x - \frac{p \cdot n}{P \cdot n}\right), \quad (3)$$

with

$$T_{ba}^i(P, p) = \frac{1}{(2\pi)^4} \int d^4 y e^{-ip \cdot y} \langle P | \bar{\psi}_b(y) \psi_a(0) | P \rangle, \quad (4)$$

where $n_\mu = (n_+, n_-, n_\perp) = (1, 0, 0_\perp)$ and P is the momentum of the hadron. The derivation of the parton evolution equations to a given order in the strong coupling, begins by expanding T_{ba}^i in perturbation theory in α_s , and evaluating the corresponding Feynman diagrams at large Q^2 and P with $x = Q^2/2P \cdot q$ fixed.

The main idea followed in Ref. 6 is to calculate the *change* in the parton distribution function from gluon radiation processes. In the infinite momentum frame as the scale Q^2 changes the number of partons with momentum fraction x also changes. Consider the number of partons with momentum fraction x at the scale Q^2 . On the one hand, partons with larger momentum fraction $x_1 > x$ at the scale Q^2 that emit a gluon with momentum fraction $x_1 - x$ now enter

into the “phase space cell” with fractional momentum x . On the other hand, partons that are already in the “phase space cell” with momentum fraction x at the scale Q^2 leave this “cell” by diminishing their value of x upon radiating a gluon. The former corresponds to the “gain” contribution, whereas the latter corresponds to the “loss” contribution.

Collins and Qiu showed that the real gluon emission processes gives a positive contribution to the parton distribution function and can be calculated from the real gluon diagram by introducing a cutoff Λ^2 [6]. Most importantly, by converting the virtual gluon diagrams into a real diagram which has the same form as that corresponding to real gluon emission except for having the opposite sign, they obtained an explicit cancelation of infrared divergences between diagrams and the “gain” and “loss” contributions to the parton distribution function [6]

$$\Delta q_i(x, \Lambda^2) = \Delta q_i(x, \Lambda^2)_{\text{gain}} - \Delta q_i(x, \Lambda^2)_{\text{loss}}, \quad (5)$$

with

$$\begin{aligned} \Delta q_i(x, \Lambda^2)_{\text{gain}} &= \int^{\Lambda^2} \frac{dk_{\perp}^2}{2\pi k_{\perp}^2} \alpha_s(k_{\perp}^2) \int_x^1 \frac{dx_1}{x_1} q_i(x_1, k_{\perp}^2) \gamma_{qq}\left(\frac{x}{x_1}\right), \\ \Delta q_i(x, \Lambda^2)_{\text{loss}} &= \int^{\Lambda^2} \frac{dk_{\perp}^2}{2\pi k_{\perp}^2} \alpha_s(k_{\perp}^2) q_i(x, k_{\perp}^2) \int_0^1 dz \gamma_{qq}(z), \end{aligned} \quad (6)$$

where the quark-quark splitting function $\gamma_{qq}(z)$ is given by Eq. (2). Taking the derivative with respect to $\ln \Lambda^2$ and identifying the cutoff scale Λ^2 as the scale Q^2 where the parton distribution function is probed [6], Collins and Qiu obtained the parton evolution equation for valence quarks as given by Eq. (1).

A close examination the above equations reveal the following important features in their derivation:

- (i) The quark distribution function that enters the right-hand side of the gain and loss contributions given by Eq. (6) carries the $k_{\perp}^2$ dependence. Since $k_{\perp}^2$ is integrated out to give the total change $\Delta q_i(x, \Lambda^2)$, this in turn is tantamount to an implicit resummation of the leading logarithms in Collins and Qiu’s derivation [6] (see also below).
- (ii) The strong coupling α_s is taken to be *running* with the scale of the exchanged gluon in the process. While the lowest order calculation of the relevant Feynman diagram given by Fig. 5 in Ref. 6 would be in terms of the bare gluon propagator and the bare strong coupling constant, in allowing the strong coupling to run with the scale, the exchanged gluon has been effectively “dressed” by parton loops.

We address these issues in detail by implementing the novel dynamical renormalization group approach to quantum kinetics [9]. This approach reveals in an alternative manner, the type of resummations implied by the final form of the parton evolution equation. This alternative formulation is based on Collins and Qiu’s probabilistic interpretation of the parton evolution equation as a Boltzmann kinetic equation with $\ln Q^2$ interpreted as a “time” variable and x as a “momentum” or “cell” variable in phase space. In particular, as will be shown below, the dynamical renormalization group approach to the Altarelli-Parisi-Lipatov equations not only provides and justifies an alternative and explicit resummation of the leading logarithms in the form a resummation of the leading secular terms, but also indicates unambiguously the running of the strong coupling in Collins and Qiu’s derivation is crucial to violations of scaling.

The derivation of the kinetic equation for the *single-particle distribution functions* in phase space via the dynamical renormalization group [9] begins by considering that the distribution function is determined at some initial time t_0 at which the initial state of the nonequilibrium system is specified. That is, the initial condition for the time evolution is determined by the initial distribution function $n_{\mathbf{k}}(t_0)$ for particles of momentum $\mathbf{k}$. Time evolution through the interaction, changes the number of particles in a given phase space cell, say, labeled by momentum $\mathbf{k}$. The evolved distribution function at a later time $t = t_0 + \Delta t$ is calculated in perturbation theory and can be written as $n_{\mathbf{k}}(t) = n_{\mathbf{k}}(t_0) + \Delta n_{\mathbf{k}}[n_{\mathbf{k}}(t_0), \Delta t]$. The change in the distribution function depends on the distribution function at the initial time $n_{\mathbf{k}}(t_0)$ because the perturbative expansion is carried out with respect to the initial state [9]; the dependence on Δt is determined by the transition probability which is obtained by using time-dependent perturbation theory. For Δt much larger than any microscopic time scales, using Fermi’s golden rule one finds that the transition probability is proportional to Δt so that $\Delta n_{\mathbf{k}}[n_{\mathbf{k}}(t_0), \Delta t] = C[n_{\mathbf{k}}(t_0)] \Delta t$, where the collision term $C[n_{\mathbf{k}}(t_0)]$ depends on the distribution functions at t_0 . Hence, one obtains

$$\frac{\Delta n_{\mathbf{k}}(t)}{\Delta t} = C[n_{\mathbf{k}}(t_0)], \quad (7)$$

which is *almost* the form of the Boltzmann equation, were it not for the fact that the term on the right-hand side depends on the distribution function at the initial time. This in turn implies that the change in the distribution

function at long times is proportional to Δt . Thus perturbation theory features *secular terms*, i.e., terms that grow in time and would lead to the breakdown of the perturbative expansion.

The Boltzmann equation is obtained after a *resummation* directly in real time via the dynamical renormalization group [9]. The resummation implied by the dynamical renormalization group can be summarized as follows: Consider evolving in time from t_0 to $t_0 + \Delta t$ with Δt much larger than a typical microscopic time so that Fermi's golden rule can be applied, but not too long so that perturbation theory is still valid during this time interval. This is exactly the assumption of the separation of time scales in a kinetic description. At the end of this interval the new distribution function is calculated as $n_{\mathbf{k}}(t_0 + \Delta t) = n_{\mathbf{k}}(t_0) + C[n_{\mathbf{k}}(t_0)] \Delta t$. Now use this new distribution function as the initial value at $t_0 + \Delta t$ to evolve further another time step, i.e., the distribution function is “reset” to the new value after each evolution in a time step. This resummation procedure can be implemented consistently via the dynamical renormalization group [9] and leads to the Boltzmann kinetic equation

$$\frac{d}{dt} n_{\mathbf{k}}(t) = C[n_{\mathbf{k}}(t)], \quad (8)$$

whose solution is found to be given by

$$n_{\mathbf{k}}(t) = n_{\mathbf{k}}(t_0) + \Delta n_{\mathbf{k}}(t), \quad \Delta n_{\mathbf{k}}(t) = \int_{t_0}^t dt' C[n_{\mathbf{k}}(t')]. \quad (9)$$

A comparison of Eq. (9) and the gain and loss contributions to the change of the parton distribution given by Eq. (6) reveals clearly that a resummation is implicitly invoked in Collins and Qiu's derivation.

We now implement a resummation via the dynamical renormalization group to obtain the evolution equation for the parton distribution function in the same manner. Following Ref. 6 we now focus on the evolution equations for the distribution function of valence quarks resulting from gluon emission. Those for the sea quarks can be obtained by a straightforward generalization once the relevant Feynman diagrams have been computed.

Consider that the parton distribution $q_i(x, Q_0^2)$ at a scale Q_0^2 is given and that the evolution from the scale Q_0^2 to a larger scale Q^2 results in a change of the parton distribution as a consequence of parton radiation. These changes are calculated via a perturbative expansion in the strong coupling constant α_s by following exactly the steps summarized above and described in detail in Ref. 6. The freedom to choose the factorization and renormalization scale permits to identify the upper momentum cutoff Λ^2 as Q^2 in the loop integrals leading to $\Delta q_i(x, Q^2)$ [6]. Taking the lower limit to be the scale Q_0^2 and the distribution function that enters in the diagrams at the scale Q_0^2 consistent with the leading logarithmic approximation, we obtain

$$\begin{aligned} q_i(x, Q^2) &= q_i(x, Q_0^2) + \Delta q_i(x, Q^2; Q_0^2), \\ \Delta q_i(x, Q^2; Q_0^2) &\equiv \Delta q_i(x, Q^2; Q_0^2)_{\text{gain}} - \Delta q_i(x, Q^2; Q_0^2)_{\text{loss}}, \end{aligned} \quad (10)$$

where

$$\begin{aligned} \Delta q_i(x, Q^2; Q_0^2)_{\text{gain}} &= \int_{Q_0^2}^{Q^2} \frac{dk_{\perp}^2}{2\pi k_{\perp}^2} \alpha_s(k_{\perp}^2) \int_x^1 \frac{dx_1}{x_1} q_i(x_1, Q_0^2) \gamma_{qq} \left(\frac{x}{x_1} \right), \\ \Delta q_i(x, Q^2; Q_0^2)_{\text{loss}} &= \int_{Q_0^2}^{Q^2} \frac{dk_{\perp}^2}{2\pi k_{\perp}^2} \alpha_s(k_{\perp}^2) q_i(x, Q_0^2) \int_0^1 dz \gamma_{qq}(z), \\ \alpha_s(k_{\perp}^2) &= \frac{\alpha_0}{\ln(k_{\perp}^2/\Lambda_{\text{QCD}}^2)}, \quad \alpha_0 = \frac{12\pi}{33 - 2N_f}, \end{aligned} \quad (11)$$

and N_f is the number of flavors.

We emphasize an important aspect of the above equations that is *different* from those obtained by Collins and Qiu: the parton distribution function enters at the *initial scale* Q_0^2 . Thus in the “update” equation (10) with the gain and loss contributions given by Eq. (11) the parton distribution function is only convoluted in the variable x but not in $k_{\perp}^2$. This aspect in turn allows to obtain the characteristic scales of parton evolution already at the perturbative level. The dependence of the updated parton evolution on the *initial* parton distribution $q_i(x, Q_0^2)$ is reminiscent of the derivation of the Boltzmann kinetic equation from the time evolution of a given *initial* state of a nonequilibrium system outlined above.

Although, following Collins and Qiu, we have included the running of the strong coupling in the gain and loss terms, we now study in detail the justification for this choice. Consider first the above expression in strict perturbation theory by taking the strong coupling to be a constant $\alpha_s(Q_0)$ fixed at the initial scale Q_0 . This corresponds to considering a bare exchanged gluon in the corresponding Feynman diagram. Performing the $k_{\perp}^2$ integral, one obtains

$$\Delta q_i(x, Q^2; Q_0^2) \propto \alpha_s(Q_0^2) \ln(Q^2/Q_0^2). \quad (12)$$

In the interpretation of $\ln Q^2$ as the “time” variable t , this in turn translates into the statement

$$\Delta q_i(x, t; t_0) \propto \alpha_s(t_0) (t - t_0), \quad (13)$$

which is akin to the expression for the *real-time* evolution of a single-particle distribution function calculated in strict perturbation theory by using Fermi’s golden rule as described above.

In the dynamical renormalization group approach [9, 11] these type of terms (divergences in the long-time limit) are identified as *secular divergences* in the perturbative expansion which signal the breakdown of perturbation theory and can be resummed systematically via the dynamical renormalization group.

Equation (12) reveals that the parton distribution function evolves on “time” scales (not to be confused with the real time) *comparable* to the scale of the running of the strong coupling constant. Therefore during the “time” scale of evolution of the parton distribution the coupling actually changes substantially and its “time” evolution cannot be neglected. This is to say that keeping the strong coupling as a constant (determined at the initial scale) does *not* allow a clear separation of “time” scales between the “kinetic” scales characteristic to parton evolution and the “microscopic” scale determined by the strong coupling since they are of the same order in the logarithmic approximation.

Including the running of the strong coupling in the loop momentum integral is tantamount to including the vacuum polarization in the exchanged gluon and leads to

$$\Delta q_i(x, Q^2; Q_0^2) \propto \alpha_0 \ln \left[\frac{\ln(Q^2/\Lambda_{\text{QCD}}^2)}{\ln(Q_0^2/\Lambda_{\text{QCD}}^2)} \right], \quad (14)$$

or, in terms of the “time” variable t ,

$$\Delta q_i(x, t; t_0) \propto \alpha_0 \ln(t/t_0). \quad (15)$$

Therefore, including the running of the coupling in the evolution kernel leads to a *slower* evolution of the parton distribution function. This is clearly a consequence of the fact that during the “time” scale of evolution of the distribution function the strong coupling changes substantially, becoming smaller and therefore in turn slowing down the change in the parton distribution.

This is the important issue of “time” scales that emerges in the kinetic description. If the coupling varies on “time” scales *longer* than that of parton evolution, then it can be kept constant during the evolution since there is a wide separation of “time” scales. However, if the coupling varies substantially during the “time” scale of parton evolution, then the running of the coupling must be included in the “update” equation of the parton distribution function. Hence, this formulation based on the “update” equation (10) with the gain and loss contributions given by Eq. (11) in terms of the parton distribution function at the initial scales allows to clarify unambiguously whether or not to include the running of the strong coupling by comparing the “time” scale of parton evolution and that of the running coupling. The criterion is as follows: If the scales are comparable or the parton distribution evolves *slower*, then the running of the coupling must be included; if the parton evolution evolves much faster, then the coupling can be kept fixed.

In the case of evolution of the parton distribution functions in QCD, the running of the strong coupling must be included in the evolution kernel of the “update” equation. A similar situation in which the evolution kernel depends explicitly on time has been found in studying quantum kinetic of charged particles in a hot QED (QCD) plasma as a result of infrared divergences associated with the emission and absorption of quasistatic, soft magnetic photons (gluons) [9, 12]. The similarity between these cases is tantalizing.

Furthermore, the above “update” equation (10) with the gain and loss contributions given by Eq. (11) implies that for fixed x the change in the parton distribution function due to a change in momentum scale only depends on the parton distribution at the initial scale Q_0^2 , i.e., the momentum scale at which the gluon is emitted, rather than on the parton distribution at intermediate scales between Q_0^2 and Q^2 . We want to stress once again that this result is consistent with the leading logarithmic approximation where gluon emission at each vertex is independent of the others and depends only on the momentum scale at which the gluon is emitted. It is this feature of the “update” equation that establishes a correspondence between the leading logarithmic approximation in parton evolution and the Markovian approximation in a kinetic description [10].

The logarithmic divergences for large Q^2 either in the form of Eq. (12) or (14) clearly signals the breakdown of the perturbative expansion and the necessity for resummation. These divergences are akin to the secular divergences that appear in time-dependent perturbation theory, in the derivation of kinetic equations [9], as well as in the asymptotic analysis of differential equations [11]. A systematic method based on the idea of renormalization group has been developed in Ref. 11 to obtain asymptotically convergent solutions of differential equations whose perturbative solution features secular terms. This renormalization group method, which we now follow, has been recently generalized to the realm of nonequilibrium quantum field theory and quantum kinetics [9].

The main idea of the renormalization group method, as described above in obtaining the Boltzmann equation, is to evolve the parton distribution function from an initial scale Q_0^2 in perturbation theory up to a larger scale $Q_0^2 + \Delta Q^2$. The change in scale ΔQ^2 is small enough so that the change in the parton distribution is perturbatively small and hence the perturbative expansion is reliable. The change in the parton distribution is absorbed into the updated parton distribution at the scale $Q_0^2 + \Delta Q^2$, which is then used for the next evolution in scale, and so on. This procedure is implemented systematically by introducing a “renormalization constant” of the *initial* parton distribution $\mathcal{Z}(x, \tau^2, Q_0^2)$ at an intermediate but otherwise arbitrary scale τ^2 , i.e.,

$$\begin{aligned} q_i(x, Q_0^2) &= \mathcal{Z}(x, \tau^2, Q_0^2) q_i(x, \tau^2), \\ \mathcal{Z}(x, \tau^2, Q_0^2) &= 1 + z^{(1)}(x, \tau^2, Q_0^2) + z^{(2)}(x, \tau^2, Q_0^2) + \dots \end{aligned} \quad (16)$$

where $q_i(x, \tau^2)$ is the “renormalized” parton distribution at scale τ^2 and $z^{(1)}(x, \tau^2, Q_0^2) \sim \mathcal{O}(\alpha_s)$, $z^{(2)}(x, \tau^2, Q_0^2) \sim \mathcal{O}(\alpha_s^2)$, etc. Substituting Eq. (16) into Eq. (10) and keeping terms only up to $\mathcal{O}(\alpha_s)$, one obtains

$$\begin{aligned} q_i(x, Q^2) &= q_i(x, \tau^2) [1 + z^{(1)}(x, \tau^2, Q_0^2)] + \int_{Q_0^2}^{Q^2} \frac{dk_\perp^2}{2\pi k_\perp^2} \alpha_s(k_\perp^2) \int_x^1 \frac{dx_1}{x_1} q_i(x_1, \tau^2) \\ &\quad - \int_{Q_0^2}^{Q^2} \frac{dk_\perp^2}{2\pi k_\perp^2} \alpha_s(k_\perp^2) q_i(x, \tau^2) \int_0^1 dz \gamma_{qq}(z) + \mathcal{O}(\alpha_s^2). \end{aligned} \quad (17)$$

The coefficients $z^{(i)}(x, \tau^2, Q_0^2)$ are chosen to cancel exactly the secular (logarithmic) terms at the scale τ order by order in the perturbative expansion. Thus, to $\mathcal{O}(\alpha_s)$, one obtains

$$\begin{aligned} z^{(1)}(x, \tau^2, Q_0^2) &= -\frac{1}{q_i(x, \tau^2)} \left[\int_{Q_0^2}^{\tau^2} \frac{dk_\perp^2}{2\pi k_\perp^2} \alpha_s(k_\perp^2) \int_x^1 \frac{dx_1}{x_1} q_i(x_1, \tau^2) \gamma_{qq}\left(\frac{x}{x_1}\right) \right. \\ &\quad \left. - \int_{Q_0^2}^{\tau^2} \frac{dk_\perp^2}{2\pi k_\perp^2} \alpha_s(k_\perp^2) q_i(x, \tau^2) \int_0^1 dz \gamma_{qq}(z) \right]. \end{aligned} \quad (18)$$

Upon substituting this choice for $z^{(1)}(x, \tau^2, Q_0^2)$ back into Eq. (17), we arrive at

$$\begin{aligned} q_i(x, Q^2) &= q_i(x, \tau^2) + \int_{\tau^2}^{Q^2} \frac{dk_\perp^2}{2\pi k_\perp^2} \alpha_s(k_\perp^2) \int_x^1 \frac{dx_1}{x_1} q_i(x_1, \tau^2) \gamma_{qq}\left(\frac{x}{x_1}\right) \\ &\quad - \int_{\tau^2}^{Q^2} \frac{dk_\perp^2}{2\pi k_\perp^2} \alpha_s(k_\perp^2) q_i(x, \tau^2) \int_0^1 dz \gamma_{qq}(z) + \mathcal{O}(\alpha_s^2), \end{aligned} \quad (19)$$

which can be improved by choosing the arbitrary renormalization scale τ^2 close to the scale Q^2 . This renormalization procedure embodies the original idea behind the Altarelli-Parisi-Lipatov equation that the change in the parton distribution due to a change in the scale can be absorbed into a redefinition, indeed a renormalization, of the parton distribution.

Since the scale τ^2 is arbitrary, a change in the renormalization scale is compensated by the corresponding change in the “renormalized” parton distribution so that $q_i(x, Q^2)$ is independent of the renormalization scale τ^2 . This in turn leads to the *dynamical renormalization group equation* $\tau^2 dq_i(x, Q^2)/d\tau^2 = 0$, which to lowest order in α_s is given by

$$\tau^2 \frac{d}{d\tau^2} q_i(x, \tau^2) = \frac{\alpha_s(\tau^2)}{2\pi} \left[\int_x^1 \frac{dx_1}{x_1} q_i(x_1, \tau^2) \gamma_{qq}\left(\frac{x}{x_1}\right) - q_i(x, \tau^2) \int_0^1 dz \gamma_{qq}(z) \right]. \quad (20)$$

Upon identifying the renormalization scale τ^2 as the scale Q^2 , one finds immediately that Eq. (20) is exactly the Altarelli-Parisi-Lipatov equation derived by Collins and Qiu. Consequently, we find that while the dynamical renormalization group leads to a kinetic equation that resums the *leading secular terms* in the perturbative expansion in nonequilibrium quantum statistical mechanics, the Altarelli-Parisi-Lipatov equation resums the *leading logarithms*, which as argued above are identified with the secular terms in a kinetic description.

The solutions of Eq. (20) can be found by taking moments of the parton distribution and quark-quark splitting functions [6]

$$\begin{aligned} F_i^n(Q^2) &= \int_0^1 dx x^{n-1} q_i(x, Q^2), \\ \gamma_{qq}^n &= \int_0^1 dz (z^{n-1} - 1) \gamma_{qq}(z). \end{aligned} \quad (21)$$

After some algebra, one finds

$$Q^2 \frac{d}{dQ^2} F_i^n(Q^2) = -\frac{\alpha_s(Q^2)}{2\pi} \gamma_{qq}^n F_i^n(Q^2), \quad (22)$$

where

$$\gamma_{qq}^n = \frac{N_c^2 - 1}{2N_c} \left[\frac{1}{2} - \frac{1}{n(n+1)} + 2 \sum_{l=2}^n \frac{1}{l} \right]. \quad (23)$$

With the running strong coupling given by Eq. (11), the solution is found to be given by

$$F_i^n(Q^2) = F_i^n(Q_0^2) \left[\frac{\ln(Q^2/\Lambda_{\text{QCD}}^2)}{\ln(Q_0^2/\Lambda_{\text{QCD}}^2)} \right]^{-\alpha_0 \gamma_{qq}^n / 2\pi}. \quad (24)$$

The mild violation of scaling, a logarithm raised to a power, instead of a pure power law with an anomalous dimension is a consequence of the running of the strong coupling inside the evolution kernel. Within the kinetic interpretation of parton evolution afforded by the dynamical renormalization group, this is a consequence of the fact recognized above at the perturbative level that the coupling runs on scales *shorter* than that of the distribution function and hence the full running coupling *must* be included in the kernel [see the discussion after Eq. (11)].

In conclusion, we have given an alternative derivation of the Altarelli-Parisi-Lipatov equation from the point of view of the dynamical renormalization group. While the final equation of evolution for the parton distribution function is the same as that obtained by Collins and Qiu [6], we believe that the derivation via the dynamical renormalization group provides further insight into the nature of the approximation and resummation. This approach helps clarify several important issues in the original derivation of Collins and Qiu:

- (i) At the level of perturbation theory, a detailed analysis of the “update” equation (10) with the gain and loss contributions given by Eq. (11) reveals that a consistent treatment of the parton evolution to leading logarithmic order should only involve the parton distribution at the initial scale. In turn this permits to understand unambiguously whether the strong coupling should be allowed to run during the scale evolution. This analysis is similar to the issue of separation of time scales in a kinetic description of nonequilibrium phenomena. It is precisely this analysis that ultimately leads to the correct violations of scaling [5], as a power of logarithms in Eq. (24), a consequence of the running of the coupling in the evolution kernel.
- (ii) The dynamical renormalization group manifestly and systematically treats the $\ln Q^2$ corrections that arises in the perturbative expansion as a renormalization of the parton distribution functions and establishes a direct correspondence between the leading logarithmic divergences arising from soft gluon emission and the leading secular divergences in time in the derivation of the Boltzmann equation. A common feature of such divergences in the parton evolution equations and in quantum kinetics is that they signal the breakdown of the perturbative approach to the evolution dynamics for large-momentum or long-time scales respectively. This breakdown of the perturbative expansion requires a resummations which is consistently recognized as a renormalization of the distribution functions.
- (iii) The dynamical renormalization group approach reveals that the leading logarithmic approximation in which the parton evolution equation only depends on the parton distribution at the momentum scale at which it is probed, corresponds to the Markovian approximation in a kinetic description.

We hope that the experience with non-Markovian kinetics in nonequilibrium statistical physics can pave the way, via the dynamical renormalization group, to understand how to consistently treat next-to or next-next-to leading logarithmic corrections. Beyond the leading logarithmic approximation lack of angular ordering of gluon emission will likely result in convolutions both in the variable x and in the variable Q^2 , i.e., a non-Markovian parton evolution in Q^2 within the kinetic interpretation. Work on these aspects is currently underway.

Acknowledgements. D.B. and H.L.Y. thank J. Qiu for illuminating discussions. The work of D.B. and S.-Y.W. was supported in part by the US NSF under grants PHY-9988720, NSF-INT-9815064 and NSF-INT-9905954. S.-Y.W. would like to thank the Andrew Mellon Foundation for partial support. D.-S.L. was supported by the ROC NSC through grant NSC-90-2112-M-259-010 and NSC-90-2112-M-259-011. H.-L.Y. would like to thank LPTHE, Université Pierre et Marie Curie (Paris VI) et Denis Diderot (Paris VII) and the Department of Physics and Astronomy, University

of Pittsburgh for their hospitality. H.-L.Y. was partially supported by the ROC NSC under grant NSC-90-2112-M-001-049.

-
- [1] H. Georgi and H.D. Politzer, Phys. Rev. D **9**, 416 (1974); D.J. Gross and F. Wilczek, 980 (1974).
 - [2] L.N. Lipatov, Yad. Fiz. **20**, 181 (1974) [Sov. J. Nucl. Phys. **20**, 94 (1975)].
 - [3] G. Altarelli and G. Parisi, Nucl. Phys. B **126**, 298 (1977); G. Altarelli, Phys. Rep. **81**, 1 (1982).
 - [4] For a pedagogical introduction to the Altarelli-Parisi-Lipatov equations, see M.E. Peskin and D.V. Schroeder, *An Introduction to Quantum Field Theory* (Addison Wesley, New York, 1995).
 - [5] CTEQ Collaboration, G. Sterman *et al.*, Rev. Mod. Phys. **67**, 157 (1995).
 - [6] J.C. Collins and J. Qiu, Phys. Rev. D **39**, 1398 (1989).
 - [7] C.F. Weizsäcker, Z. Phys. **88**, 612 (1934); E.J. Williams, Phys. Rev. **45**, 729 (1934).
 - [8] L. Durand and W. Putikka, Phys. Rev. D **36**, 2840, (1987).
 - [9] D. Boyanovsky, H. J. de Vega, R. Holman and M. Simionato, Phys. Rev. D **60**, 065003 (1999); D. Boyanovsky and H.J. de Vega, **59**, 105019 (1999); D. Boyanovsky, H.J. de Vega, and S.-Y. Wang, **61**, 065006 (2000); S.-Y. Wang, D. Boyanovsky, H.J. de Vega, and D.-S. Lee, **62**, 105026 (2000).
 - [10] For instance, see R. Balescu, *Equilibrium and Nonequilibrium Statistical Mechanics* (Wiley, New York, 1975).
 - [11] L.-Y. Chen, N. Goldenfeld, and Y. Oono, Phys. Rev. Lett. **73**, 1311 (1994); Phys. Rev. E **54**, 376 (1996); T. Kunihiro, Prog. Theor. Phys. **95**, 503 (1995); **97**, 179 (1997); S. Ei, K. Fujii, and T. Kunihiro, Ann. Phys. (N.Y.) **280**, 236 (2000).
 - [12] J.-P. Blaizot and E. Iancu, Phys. Rev. Lett. **76**, 3080 (1996); Phys. Rev. D **55**, 973 (1997); **56**, 7877 (1997).